



Entrez PubMed

Nucleotide

Protein

Genome

Structure

PMC

Journals

Bo

Search PubMed

for

Go

Clear

☒ Limits

Preview/Index

History

Clipboard

Details

About Entrez

Display

Abstract

Show:

20

Sort

Send to

Text

Text Version

☐ 1: Biochemistry. 1992 Mar 31;31(12):3281-8.

Related Articles, Link

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Database

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

Cubby

Related Resources

Order Documents

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

Privacy Policy

Biochemical and spectroscopic characterization of the high molecular weight cytochrome c from *Desulfovibrio vulgaris* Hildenborough expressed in *Desulfovibrio desulfuricans* G200.

Bruschi M, Bertrand P, More C, Leroy G, Bonicel J, Haladjian J, Chottard G, Pollock WB, Voordouw G.

Laboratoire de Chimie Bacterienne, Centre National de la Recherche Scientifique, Marseille, France.

The gene of high molecular weight, multiheme cytochrome c (Hmc) from the sulfate-reducing bacterium *Desulfovibrio vulgaris* Hildenborough has been overexpressed in *Desulfovibrio desulfuricans* G200. The recombinant protein has been purified. Its molecular weight (65,600), amino acid composition, and NH₂-terminal sequence were found to be identical to those of the wild-type protein. The recombinant protein has been spectroscopically characterized (optical spectrum, EPR, circular dichroism) and compared to the wild-type protein. We have found 16 hemes per molecule by iron analysis and the pyridine hemochrome test. Both high- and low-spin features were observed in the EPR spectrum. A detailed spin quantitation analysis indicates 1 or 2 high-spin hemes and 14 or 15 low-spin hemes per molecule. The redox potentials of the hemes determined by voltammetric techniques gave an average of three different values, 0, -100, and -250 mV (versus NHE), for the wild-type and the recombinant cytochrome. The low potential values are similar to the values observed for the bis(histidiny) coordinated hemes of cytochrome c₃. A comparison of the arrangement of heme binding sites and coordinated histidines in the amino acid sequences of cytochrome c₃ and Hmc has shown that the latter contains four domains, three of which are complete c₃-like domains, while the fourth represents an incomplete c₃-like domain which may contain His-Met coordinated hemes. These data are in agreement with the detailed study of the number and types of hemes reported in this paper.

PMID: 1313289 [PubMed - indexed for MEDLINE]

Display

Abstract

Show:

20

Sort

Send to

Text

[Write to the Help Desk](#)
[NCBI](#) | [NLM](#) | [NIH](#)
[Department of Health & Human Services](#)
[Freedom of Information Act](#) | [Disclaimer](#)

Oct 2 2003 18:06: